

Prenatrix™

Dietary Supplement Multivitamin (30ct)

NDC: 59088-166-54



DERMACINRx

Description:

Prenatrix™ is an orally administered prescription product that provides vitamin supplementation during and after pregnancy.

Prenatrix™ works well as postnatal vitamins to continue to provide important nutrients for your baby while breastfeeding and may be continued to be taken throughout a mother's childbearing years.

Unit Detail:

30 ct per bottle

Dosage:

One caplet daily

See reverse for complete prescribing info

Prenatrix™

Multivitamin Rx Only

Each caplet contains:

Vitamin A (as retinyl acetate).....	1500 mcg (5000 IU)
Vitamin C (as ascorbic acid).....	120 mg
Vitamin D ₃ (as cholecalciferol).....	20 mcg (800 IU)
Vitamin E (dl-alpha tocopheryl acetate).....	30 mg (30 IU)
Thiamin (as thiamine mononitrate).....	3 mg
Riboflavin (vitamin B ₂).....	3.4 mg
Niacin (as niacinamide).....	20 mg
Vitamin B ₆ (as pyridoxine hydrochloride).....	50 mg
Folate (as folic acid).....	1700 mcg DFE (1000 mcg folic acid)
Vitamin B ₁₂ (as cyanocobalamin).....	10 mcg
Choline (as choline bitartrate).....	55mg
Calcium (as calcium carbonate).....	200 mg
Iron (as ferrous fumarate).....	27 mg
Iodine (as potassium iodine).....	150 mcg
Magnesium (as magnesium oxide).....	200 mg
Zinc (as zinc oxide).....	25 mg
Selenium (as selenium amino acid chelate).....	70 mcg
Manganese (as manganese sulfate).....	2.6 mg
Chromium (as chromium polynicotinate).....	45 mcg
Molybdenum (as molybdenum amino acid chelate).....	50 mcg

Each Bottle Contains: 30 Caplets

Other Ingredients: BHT, dicalcium phosphate, croscarmellose sodium, crospovidone, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, modified food starch, pork gelatin, starch aluminium octenyl succinate, sodium ascorbate, sodium aluminum silicate, silicon dioxide, stearic acid, sucrose, Clear Coating: (hydroxypropyl methylcellulose, PEG-8).

Contraindications: This product is contraindicated in patients with known hypersensitivity to any of its ingredients; also, all iron compounds are contraindicated in patients with hemosiderosis, hemochromatosis, or hemolytic anemias. Pernicious anemia is a contraindication, as folic acid may obscure its signs and symptoms.

Warnings: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Administration of folic acid alone is improper therapy for pernicious anemia and other megaloblastic anemias in which vitamin B12 is deficient.

Indications and usage:

Prenatrix™ is used during pregnancy and during the postnatal period as prescribed by your health care provider for both lactating and non-lactating mothers, and may be continued to be taken throughout a mother's childbearing years.

Prenatrix™ may be beneficial in improving the nutritional status of women prior to conception.

For use on the order of a healthcare practitioner.

Call your doctor about side effects. To report side effects, call **PureTek Corporation** at **1-877-921-7873** or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Store at controlled room temperature 20° to 25°C (68° to 77°F). [See USP]. Protect from light and moisture and avoid excessive heat.

Do not use if bottle seal is broken.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Store at controlled room temperature 20° to 25°C (68° to 77°F). [See USP].

Protect from light and moisture and avoid excessive heat.

To report a serious adverse event or to obtain product information, contact **877-921-7873**.

***Vitamin supplement labels now list “mcg DFE of folate,” which stands for dietary folate equivalent. It’s the amount of folate your body absorbs.**

Manufactured by:
PureTek Corporation
Panorama City, CA 91402
For questions or information
call toll-free: **877-921-7873**